

1. Introduction

First order electroweak phase transition has often been discussed [1] within the context of the finite temperature effective potential of the order parameter, which is the expectation value of the Higgs field, evolving in a thermal background. Such phase transitions are highly local phenomena: critical bubbles have a finite size. However, effective potentials are averaged over all space and do not describe local fluctuations, which may affect the dynamics of nucleation and bubble growth. Fluctuations with spatial correlations comparable to the critical bubble size may be expected to be important for bubble nucleation and, therefore, we concentrate on the thermalization of an on-shell Higgs field with zero spatial momentum ($p_0 = m_H$, $\mathbf{p} = 0$). This choice reflects the fact that critical bubbles are typically much bigger than the inter particle distance $\sim 1/T$ in plasma.

Local fluctuations of the order parameter, which one might call subcritical fluctuations, are continuously created and destroyed by the self-interactions of the thermal background. A natural way to describe them is to consider a non-constant order parameter, for which one needs the complete effective action for the background field $\phi(x)$. The computation of the effective action is in general very difficult, but if one focuses on small-amplitude fluctuations, $\phi(x) = \phi_0 + b(x)$ with $b(x)$ small, one may make use of a perturbation expansion in $b(x)$, and to order b^2 the finite temperature effective action reads simply [2]

$$\Gamma[\phi_0, b] = -V_{eff}(\phi_0) + \Gamma^{(1)}(\phi_0)b(k=0) + \frac{1}{2} \int \frac{d^4k}{(2\pi)^4} b(-k) \Gamma^{(2)}(\phi_0; k) b(k) , \quad (1)$$

where the tadpole $\Gamma^{(1)} = 0$ at the high temperature minimum $\phi_0 = 0$, and $\Gamma^{(2)}$ is the two-point function for the Higgs field, which can be computed in ordinary perturbation theory.

Two-loop dispersion relations and the evolution of the Fourier modes $b(k)$ in $\lambda\phi^4$ -theories has been discussed in detail in [2]. In the present paper we investigate how fast the fluctuations thermalize in the Standard Model above, and just prior to, the electroweak phase transition. We assume the standard

particle content and one Higgs doublet. The thermalization rate γ depends on the imaginary part of the effective action $\text{Im } \Sigma \equiv -\text{Im } \Gamma^{(2)}$, via

$$\gamma = -\frac{\text{Im } \Sigma}{\omega}, \quad (2)$$

where $\omega = \omega(k)$ is the energy of a given mode. We compute the imaginary part of the two-point function at one-loop level for fermion and gauge boson corrections (at one loop there is no higgs contribution to $\text{Im } \Sigma$ in the unbroken phase). The imaginary parts are cuts of the relevant one-loop diagrams [3,4], and therefore equivalent to processes which may be called decay, inverse decay, absorption, or emission of the higgs. For massless fermions and gauge bosons these processes are kinematically forbidden at one-loop level. However, the physical external states should include thermal corrections which, to leading order in T , are the hard thermal loops. They merely change the pole structure of the propagators to account for the interactions with the thermal background. Accordingly, there appear energy thresholds, which in particular for the fermionic loops are rather complicated because of the complicated nature of the dispersion relations. Here, the one-loop case is exceptional in that at higher loops there are no thresholds.

The paper is organized as follows. Section 2 contains a discussion about the fermion dispersion relations at finite temperature and different channels, and we present a derivation of the fermionic contribution to the thermalization rate. In Section 3 we compute the gauge boson loop and estimate also the size of the two-loop corrections. In Section 4 we compare the thermalization rate to the electroweak phase transition rate, and Section 5 contains our conclusions together with a summary of our results.

2. Fermions

2.1 Preliminaries

The dispersion relation of massless chiral fermions is drastically changed by the high temperature correction [5,6]. The leading T^2 part of the fermion self-energy gives rise to two kinds of excitations of the thermal heat bath called particles and holes [7]. Both particles and holes have unusual dispersions relations. Particles have a minimum group velocity that is 1/3 of the speed of light, so that there are no particle excitations at rest, even though there is a mass gap. This phenomenon is due to the collective excitations. Holes have the same group velocity for $k = 0$ but with an opposite sign, and a minimum energy for approximately $k = 0.408 m_f$ [8]. All these features combined together make the thresholds difficult to determine. The dispersion relations for particles and holes are displayed schematically in Fig. 1.

Before studying the thresholds of the diagram in Fig. 2 we need an expression for $\text{Im } \Sigma$ including all possible channels. We may use the expression in [4] suitably modified to account for collective excitations and chirality. One difference between the right and left handed fermion propagators is thermally induced masses $m_{L,R}$ which are not equal because the right and left handed fermions are in different representations of the Standard Model gauge group. The one-loop resummed propagator in the rest frame of the heat bath reads ($P_\mu = (p_0, \mathbf{p})$ with $p = |\mathbf{p}|$)

$$S(P) = \eta \frac{\not{L}(P)}{L^2(P)} \bar{\eta} + \bar{\eta} \frac{\not{R}(P)}{R^2(P)} \eta , \quad (3)$$

where

$$\begin{aligned} \eta &= \frac{1}{2}(1 - \gamma_5) , \quad \bar{\eta} = \frac{1}{2}(1 + \gamma_5) , \\ \not{L} &= (1 + a_L(p_0, p)) P_\mu \gamma^\mu + b_L(p_0, p) \gamma^0 , \\ a_L(p_0, p) &= \frac{m_L^2}{p^2} \left(1 - \frac{p_0}{2p} \ln \left| \frac{p_0 + p}{p_0 - p} \right| \right) , \\ b_L(p_0, p) &= \frac{m_L^2}{p} \left(-\frac{p_0}{p} + \left(\frac{p_0^2}{p^2} - 1 \right) \frac{1}{2} \ln \left| \frac{p_0 + p}{p_0 - p} \right| \right) , \end{aligned} \quad (4)$$

and similarly for $\bar{R}$. The functions $a_{L,R}(p_0, p)$ and $b_{L,R}(p_0, p)$ are respectively even and odd with respect p_0 , i.e. they have the properties

$$a_{L,R}(-p_0, p) = a_{L,R}(p_0, p) , \quad b_{L,R}(-p_0, p) = -b_{L,R}(p_0, p) . \quad (5)$$

We have introduced left and right handed fermion masses in Eq. (4). For leptons one finds

$$\begin{aligned} m_{l,L}^2 &= \frac{m_Z^2 + 2m_W^2 + m_l^2 + m_{l'}^2}{2f^2} T^2, \\ m_{l,R}^2 &= \frac{m_Z^2 - m_W^2 + \frac{1}{2}m_l^2}{2f^2} T^2, \end{aligned} \quad (6)$$

where m_l and $m_{l'}$ are the $T = 0$ masses of the leptons l and l' belonging to the same $SU(2)$ doublet. For quarks one finds that

$$\begin{aligned} m_{u_i,L}^2 &= \frac{1}{6}g_s^2 T^2 + \frac{3m_W^2 + \frac{1}{9}(m_Z^2 - m_W^2) + m_{u_i}^2 + m_{d_i}^2}{8f^2} T^2, \\ m_{u_i,R}^2 &= \frac{1}{6}g_s^2 T^2 + \frac{\frac{4}{9}(m_Z^2 - m_W^2) + \frac{1}{2}m_{u_i}^2}{2f^2} T^2, \end{aligned} \quad (7)$$

and

$$\begin{aligned} m_{d_i,L}^2 &= \frac{1}{6}g_s^2 T^2 + \frac{3m_W^2 + \frac{1}{9}(m_Z^2 - m_W^2) + m_{u_i}^2 + m_{d_i}^2}{8f^2} T^2, \\ m_{d_i,R}^2 &= \frac{1}{6}g_s^2 T^2 + \frac{\frac{1}{9}(m_Z^2 - m_W^2) + \frac{1}{2}m_{d_i}^2}{2f^2} T^2, \end{aligned} \quad (8)$$

where m_{u_i} and m_{d_i} are the $T = 0$ masses of u and d type quarks of i -th generation, respectively, $m_W = 80.0$ GeV, $m_Z = 91.2$ GeV and m_t is the $T = 0$ mass of the top-quark. The higgs vacuum expectation value is $f \simeq 247$ GeV. For the strong coupling constant we adopt the value $g_s \simeq 1.2$. Note that the left handed masses of fermions belonging to same $SU(2)$ doublet are equal due to unbroken symmetry.

The imaginary part of the self-energy of the diagram in Fig. 2 is

$$\begin{aligned} \text{Im } \Sigma_f(p_0, \mathbf{p}) = & -\frac{4g_Y^2}{\sinh 2\theta_p} \int \frac{d^4 K}{(2\pi)^2} \epsilon(k_0) \epsilon(k_0 - p_0) \frac{1}{4} \sin 2\phi_k \sin 2\phi_{k-p} \\ & \times L_\mu(K) R^\mu(K - P) \eta_L \eta_R \delta(L^2(K)) \delta(R^2(K - P)) , \end{aligned} \quad (9)$$

where

$$\sin 2\phi_k = 2 \frac{e^{\beta|k_0|/2}}{e^{\beta|k_0|} + 1} , \quad (10)$$

and g_Y is the corresponding Yukawa coupling. The factors η_L and η_R are +1 and -1 for particles and holes, respectively. They arise when the energies in $L^2(K)$ and $R^2(K - P)$ are continued analytically using the time-ordered prescription $k_0 \rightarrow k_0 + i\epsilon k_0$. For a free massive propagator this would only add $i\epsilon$ in the denominator. In our case we have

$$L^2(k_0 + i\epsilon k_0, \mathbf{k}) = L^2(K) + i\epsilon k_0 \frac{\partial L^2}{\partial k_0} . \quad (11)$$

The factor that multiplies $i\epsilon$ is positive for particle solutions and negative for hole solutions which gives rise to an extra sign for the hole contribution. Below we shall see that this extra sign is crucial in order to get a positive damping rate in all channels, whereas the naive propagator $(L^2(K) + i\epsilon)^{-1}$ would lead to an instability.

2.2 Thresholds

From the analysis in [3] it follows that there are two possible decay channels for a scalar interacting with fermions with two different masses, m_1 and m_2 . In the case of a scalar at rest ($p_0 = m_H, \mathbf{p} = 0$) they are: (1) decay if $m_H > m_1 + m_2$, (2) absorption if $m_H < |m_1 - m_2|$. The inverses of these processes are also possible and they are included in the rates we compute below. Here, and in what follows, we denote by $m_H \equiv m_H(T)$ the T -dependent higgs mass.

In our case the thresholds are not determined by the $\mathbf{k} = 0$ energy but must be derived using the complete dispersion relations. A decay or absorption can take place when one of the following conditions is fulfilled

$$\begin{aligned} (1) \quad m_H &= \omega_{L,i}(\mathbf{k}) + \omega_{R,j}(\mathbf{k}) , \\ (2) \quad m_H + \omega_{L,i}(\mathbf{k}) &= \omega_{R,j}(\mathbf{k}) , \\ (3) \quad m_H + \omega_{R,j}(\mathbf{k}) &= \omega_{L,i}(\mathbf{k}) . \end{aligned} \quad (12)$$

Here $k=|\mathbf{k}|$ and for each energy we have put an index $(i, j = p, h)$ to distinguish between particles and holes. However, not all combinations of particles and holes contribute in each channel because the factor $L_\mu R^\mu$ can vanish. This turns out to be important in order to obtain a positive thermalization rate.

Let us denote by E the energy of an excitation, including a sign with reference to the arrows in Fig. 2. Then E_L and E_R have opposite signs in the decay process, and equal signs in the absorption process. From Eq. (4) we have explicitly

$$\begin{aligned} L_\mu(E_{L,i}, \mathbf{k}) R^\mu(E_{R,j}, \mathbf{k}) = \\ [E_{L,i}(1 + a_L(E_{L,i}, \mathbf{k})) + b_L(E_{L,i}, \mathbf{k})][E_{R,j}(1 + a_R(E_{R,j}, \mathbf{k})) + b_R(E_{R,j}, \mathbf{k})] \\ - k^2(1 + a_L(E_{L,i}, \mathbf{k}))(1 + a_R(E_{R,j}, \mathbf{k})) . \end{aligned} \quad (13)$$

This should be compared with the on-shell conditions for positive energy particles and holes

$$\begin{aligned} \omega_p(1 + a(\omega_p, \mathbf{k})) + b(\omega_p, \mathbf{k}) &= k(1 + a(\omega_p, \mathbf{k})) , \\ \omega_h(1 + a(\omega_h, \mathbf{k})) + b(\omega_h, \mathbf{k}) &= -k(1 + a(\omega_h, \mathbf{k})) . \end{aligned} \quad (14)$$

Negative energy solutions, where $\omega \rightarrow -\omega$ in Eq. (14), are also possible but then the on-shell conditions for particles and holes are reversed due to the properties of $a_{L,R}$ and $b_{L,R}$ in Eq. (5). Using Eq. (14) in Eq. (13) we see that when E_L and E_R have the same sign, only $(i = p, j = h)$ and $(i = h, j = p)$ give a non-zero result. When E_L and E_R have opposite signs we are left with $(i = p, j = p)$ and $(i = h, j = h)$. This selection rule can be understood physically as a consequence of the conservation of angular momentum. The chirality–helicity relation is reversed for holes [7], and because the fermions in scattering processes under consideration always have different chiralities, absorption is possible only if they have same helicities, and direct decay is possible only if they have opposite helicities.

For a Higgs particle at rest the angular integrals in Eq. (9) become trivial and the δ -functions can be used to perform the k_0 and $\mathbf{k}$ integrations. One should note that the δ -function may be rewritten as

$$\delta(L^2) = \sum_{i=p,h} W_{L,i}(\mathbf{k}) [\delta(k_0 - \omega_{L,i}(\mathbf{k})) + \delta(k_0 + \omega_{L,i}(\mathbf{k}))] , \quad (15)$$

where the energy factor $W_{L,i}$ is different from the usual $1/2\omega(k)$. In fact, we have instead

$$W_{L,i}^{-1}(k) = \left| \frac{\partial L^2}{\partial k_0} \right|_{k_0=\omega_{L,i}} , \quad (16)$$

and similarly for W_R . Combining the expression in Eq. (13) for $L_\mu R^\mu$ with the factors W_L and W_R we find the rather simple expression

$$L_\mu R^\mu W_L W_R = - \frac{2(\omega_{L,i}^2 - k^2)(\omega_{R,j}^2 - k^2)}{4m_L^2 4m_R^2} , \quad (17)$$

which will be useful later. We note that $L_\mu R^\mu W_L W_R$ is always negative.

2.3 Decay rates

Let us start with the absorption channel, which opens up when m_H is small enough. Its existence is determined by the last two conditions in Eq. (12). In this case E_L and E_R have the same sign so that only $p \rightarrow h$ and $h \rightarrow p$ processes contribute. By virtue of the dispersion relations (see Fig. 1) there is a channel for small m_H even if $m_L = m_R$, because when $\omega_h \leq \omega_p$ the higgs may be absorbed by a hole in the heat bath to produce a particle. Using Eq. (17) the imaginary part for the process $H + f_{R,h} \rightarrow f_{L,p}$ can then be found to be

$$\text{Im } \Sigma(Rh, Lp) = \frac{4g_Y^2}{\pi} \eta_{L,p} \eta_{R,h} k^2 \frac{(\omega_{L,p}^2 - k^2)(\omega_{R,h}^2 - k^2)}{4m_L^2 4m_R^2} [n_{R,h}(k) - n_{L,p}(k)] , \quad (18)$$

where k is determined by the relation

$$m_H + \omega_{R,h}(k) = \omega_{L,p}(k) , \quad (19)$$

and

$$n_{L,R} = \frac{1}{e^{\beta\omega_{L,R}} + 1} , \quad (20)$$

are the fermion distribution functions. We find that $\text{Im } \Sigma$ in Eq. (18) is negative, as it should, leading to a positive decay rate, but the reason is different from the case of ordinary massive fermions [3]. Here, the matrix element $L_\mu R^\mu$ is negative in all channels, in contrast to the case in [3] where the signs are different for decay

and absorption. On the other hand, we have an extra factor of $\eta_{L,i}\eta_{R,j}$ which then helps to give the correct sign. We note that this works because only $p \rightarrow h$ and $h \rightarrow p$ processes contribute in this channel. To obtain the complete fermionic $\text{Im } \Sigma$ for absorptions, the contribution from the process $H + f_{L,h} \rightarrow f_{R,p}$ should be added to Eq. (18) so that $\text{Im } \Sigma_{abs} = \text{Im } \Sigma(Rh, Lp) + \text{Im } \Sigma(Lh, Rp)$. In the minimal Standard Model the difference between m_L and m_R is so small compared to the particle-hole difference that we could safely approximate $m_L = m_R$, which would simplify the expression for $\text{Im } \Sigma_{abs}$. (See, however, the discussion below when the mass difference is not neglected.)

From Fig. 1 we see that the threshold is at a finite k_1 and two solutions around $k = k_1$ exist for m_H slightly below the threshold. When m_H decreases one solution goes to zero, if we neglect the mass difference, and therefore its contribution to $\text{Im } \Sigma$ goes to zero. The other solution goes to infinity but the residue of the hole propagator, which is proportional to $\omega_h^2 - k^2$, goes exponentially to zero and thus $\text{Im } \Sigma_{abs} \rightarrow 0$ when $m_H \rightarrow 0$.

If we do not neglect the mass difference in the absorption channel the threshold structure becomes more complicated. Let us assume that $m_L > m_R$. Then in the $H + f_{R,h} \rightarrow f_{L,p}$ channel (Fig. 1a) one of the solutions disappears when $m_H < m_L - m_R$. In the $H + f_{L,h} \rightarrow f_{R,p}$ channel (Fig. 1b) the dispersion curves cross at some point k_2 . The threshold is determined by the largest difference to the right of the crossing point (k_1 in Fig. 1b). As m_H decreases there appear two solutions to the right. In the limit $m_H \rightarrow 0$, the large k solution disappears since the hole residue goes exponentially to zero. We are then left with a solution which for a small mass difference is located at $k_2 = \frac{3}{2}(m_L - m_R)$. When $m_H \rightarrow 0$ the thermalization rate can be found from Eq. (18) to be

$$\gamma_F/T = \frac{9g_Y^2}{16\pi} \left(\frac{(m_R + m_L)^2}{4m_R m_L} \right)^2 \left(\frac{m_R - m_L}{T} \right)^2 \frac{1}{1 + \cosh[\beta(m_R + m_L)]} . \quad (21)$$

There is also a third absorption channel, $H + f_{R,p} \rightarrow f_{L,h}$, which opens for $m_H < m_L - m_R$ and k is located between 0 and k_2 . The thermalization rate in the limit $m_H \rightarrow 0$ is again given by Eq. (21).

The direct decay channel is simpler. The signs of E_L and E_R are opposite, and the higgs decays either into a pair of particles or into a pair of holes. Therefore, the product $\eta_L \eta_R$ is +1, leading again to a positive decay rate. Summing over both decay channels we find

$$\text{Im } \Sigma_{dec} = -\frac{4g^2}{\pi} \sum_{i=p,h} k^2 \frac{(\omega_{L,i}^2 - k^2)(\omega_{R,i}^2 - k^2)}{4m_L^2 4m_R^2} (1 - n_{L,i} - n_{R,i}) , \quad (22)$$

and k is in this case determined by

$$m_H = \omega_{L,i}(k) + \omega_{R,i}(k) . \quad (23)$$

The ordinary $T = 0$ contribution is included in this expression. The threshold for holes is lower than for particles since the minimum energy for holes is about $0.928 m_f$ [8]. There are also two solutions in this channel when $1.865 m_f < m_H < 2 m_f$. As m_H increases the corresponding k increases and the hole residue goes exponentially to zero so that the particle channel dominates as soon as m_H is above the particle threshold.

The full fermionic contribution is thus finally given by $\text{Im } \Sigma_f = \text{Im } \Sigma_{abs} + \text{Im } \Sigma_{dec}$. In practise the direct decay channel is in many cases irrelevant, because its existence would imply that $m_H(T) \gtrsim T$. At least in perturbation theory m_H is expected to be less than T in the high temperature limit. We shall return to $\text{Im } \Sigma_f$ in Section 4, where we also plot it for the top quark.

3. Gauge bosons

3.1 One-loop case

At $T = 0$ the imaginary part of the diagram of Fig. 3 is zero for an on-shell massive Higgs particle and a massless gauge boson for kinematical reasons. At finite temperature thermal masses are induced for both the Higgs and the gauge boson so that an absorption channel opens when $2 m_H(T) < m_W(T)$. In particular,

this condition may be fulfilled close to the critical temperature of a second order, or a weakly first order, phase transition where the higgs mass becomes small.

The self-energy of a gauge boson is a rather complicated function of the momentum [5,6]. We may, however, still use the standard expression for the imaginary part of the self-energy, which can be found in e.g. [4], if we take into account the fact that the higgs couples to gauge bosons through a derivative coupling, and replace the δ -functions with the ones containing the thermal on-shell condition. With the high T expression for the gauge boson propagator* [5,6] we get ($K_\mu = (k_0, \mathbf{k})$, $k = |\mathbf{k}|$)

$$\begin{aligned} \text{Im } \Sigma_{gb}(p_0, \mathbf{p}) = & \\ & - \frac{e^2}{\sinh 2\theta_p} \int \frac{d^4 K}{(2\pi)^2} \frac{1}{4} \sinh 2\theta_k \sinh 2\theta_{p-k} \delta((\mathbf{p} - \mathbf{k})^2 - m_H^2(T)) \\ & \times \left\{ 4 \left(\frac{(\mathbf{p} \cdot \mathbf{k})^2}{k^2} - \mathbf{p}^2 \right) \delta(K^2 - \pi_T) + \frac{4}{K^2 k^2} (p_0 k^2 - k_0 \mathbf{p} \cdot \mathbf{k})^2 \delta(K^2 - \pi_L) \right\} , \end{aligned} \quad (24)$$

where

$$\sinh 2\theta_p = 2 \frac{e^{\beta|p_0|/2}}{e^{\beta|p_0|} - 1} . \quad (25)$$

and

$$e^2 = \begin{cases} \frac{3}{4}g^2 & \text{for } SU(2) , \\ g'^2 & \text{for } U(1) . \end{cases} \quad (26)$$

The gauge boson self-energy has a transverse and a longitudinal part, which are given by

$$\pi_T(k_0, \mathbf{k}) = \frac{g^2 T^2}{6} \left((n_W + 1) \left[\frac{k_0^2}{k^2} + \left(1 - \frac{k_0^2}{k^2}\right) \frac{k_0}{2k} \ln \left| \frac{k_0 + k}{k_0 - k} \right| \right] - 2 \right) , \quad (27)$$

$$\pi_L(k_0, \mathbf{k}) = \frac{(n_W + 1)g^2 T^2}{3} \left(1 - \frac{k_0^2}{k^2}\right) \left[1 - \frac{k_0}{2k} \ln \left| \frac{k_0 + k}{k_0 - k} \right| \right] .$$

These self-energies include the contributions from gauge boson, fermion and scalar loops. We use the notation $n_W = (N_f + 4N)/4$, where $N_f = 2 \times (\# \text{ of generations})$

* We are using the Feynman gauge throughout the paper.

and $N = 2$ for $SU(2)$ gauge fields. For $U(1)$ gauge field g is replaced by g' and n_W by $n_Y = \frac{19}{36}N_f$. We define also the longitudinal and transverse gauge boson masses by $M_L^2 = \pi_L(\omega, \mathbf{k} = 0)$ and $M_T^2 = \pi_T(\omega, \mathbf{k} = 0)$, respectively.

At one loop the high T expression for the higgs mass in the unbroken phase is given by

$$m_H^2(T) = -\frac{m_H^2(0)}{2} + \frac{6m_W^2 + 3m_Z^2 + 6m_t^2 + m_H^2(0)}{12f^2} T^2, \quad (28)$$

where $m_H(0)$ is the zero temperature higgs mass. The longitudinal masses of $SU(2)$ and $U(1)$ gauge bosons are given by

$$\begin{aligned} M_{L,SU(2)}^2 &= \frac{n_W + 1}{9} \frac{4m_W^2}{f^2} T^2, \\ M_{L,U(1)}^2 &= \frac{n_Y + 1}{9} \frac{4(m_Z^2 - m_W^2)}{f^2} T^2, \end{aligned} \quad (29)$$

while the transverse masses are

$$\begin{aligned} M_{T,SU(2)}^2 &= \frac{n_W - 2}{9} \frac{4m_W^2}{f^2} T^2, \\ M_{T,U(1)}^2 &= \frac{n_Y - 2}{9} \frac{4(m_Z^2 - m_W^2)}{f^2} T^2. \end{aligned} \quad (30)$$

It is the scalar loop that gives rise to the difference between the longitudinal and transverse masses.

For a Higgs particle at rest ($p_0 = m_H, \mathbf{p} = 0$) only the longitudinal part contributes and the angular integral becomes again trivial. We obtain, as in Section 2, an energy factor from the residue at the pole, given by

$$\widetilde{W}_L^{-1}(\mathbf{k}) = \left| \frac{\partial}{\partial k_0} \left(K^2 - \pi_L(k_0, \mathbf{k}) \right) \right|_{k_0 = \omega_L(\mathbf{k})} = \frac{M_L^2 - \pi_L(\omega_L(\mathbf{k}), \mathbf{k})}{\omega_L(\mathbf{k})}. \quad (31)$$

The absorption channel opens up when the higgs mass is so small that it can annihilate with a higgs from the heat bath to produce a gauge boson. In that channel $\text{Im } \Sigma$ is given by

$$\text{Im } \Sigma_{gb} = -\frac{e^2}{\pi} \frac{\widetilde{W}_L(\mathbf{k})}{\pi_L(\mathbf{k}) \sqrt{k^2 + m_H^2(T)}} m_H^2(T) k^4 [n_H(\mathbf{k}) - n_L(\mathbf{k})], \quad (32)$$

where k is determined by solving the on-shell condition of the gauge boson relation

$$\omega_L^2(k) = k^2 + \pi_L(\omega_L(k), k) , \quad (33)$$

and the energy conservation condition

$$m_H(T) + \sqrt{k^2 + m_H^2(T)} = \sqrt{k^2 + \pi_L(\omega_L(k), k)} . \quad (34)$$

The thermal distribution functions are

$$n(k)_{H,L} = \frac{1}{e^{\beta\omega_{H,L}} - 1} , \quad (35)$$

with

$$\omega_H = \sqrt{k^2 + m_H^2(T)} , \quad \omega_L = \sqrt{k^2 + \pi_L} . \quad (36)$$

Equation (32) expresses the imaginary part for any component of the Higgs doublet; they are all equal because of the unbroken gauge symmetry.

We are mainly interested in the thermalization rate close to the phase transition where $m_H(T)$ is expected to be small. In the limit $m_H \rightarrow 0$ the thermalization rate due to $SU(2)$ gauge bosons is given by

$$\begin{aligned} \gamma_{gb}/T = \frac{3g^2}{8\pi} \left(\frac{T}{M_L} \right)^2 & \left(\frac{(n_W + 1)g^2}{6} \ln \frac{T}{m_H} \right)^{3/2} \\ & \frac{m_H}{T} \exp \left[- \left(\frac{(n_W + 1)g^2}{6} \ln \frac{T}{m_H} \right)^{1/2} \right] . \end{aligned} \quad (37)$$

This result can be found by using the asymptotic formula [5] $\pi_L = 4k^2 \exp(-6k^2/(n_W + 1)g^2T^2)$ which is valid for $3k/\sqrt{n_W + 1}gT \gg 1$, together with Eqs. (33) and (34). One solves k iteratively as a function of m_H , whence one obtains Eq. (37) from Eq. (32), and the expansion is valid when $m_H \lesssim 10^{-3}T$, corresponding to $\gamma_{gb}/T \lesssim 1.9 \times 10^{-4}$.

3.2 Two-loop case

At two-loop level the variety of relevant processes is much larger than at one loop, leading to tens of different diagrams. We now estimate the order of

magnitude of two-loop processes using the two representative graphs given in Fig. 4. The purely scalar "rising sun" diagram of Fig. 4a has been calculated in [2]. The imaginary part of the two-point function for an on-shell complex higgs with a vanishing external three momentum is, in the leading high T approximation,

$$\text{Im}\Sigma_H = -\frac{\lambda^2}{256\pi}T^2. \quad (38)$$

We may use this result to estimate the imaginary part of Fig. 4b which then reads

$$\text{Im}\Sigma = -\frac{3g^4}{256\pi}T^2 \quad (39)$$

for $SU(2)$ gauge bosons. The $U(1)$ gauge boson has a similar expression but with $3g^4$ replaced by $4(g')^4$. In the derivation of Eq. (39) the mass difference of gauge bosons and Higgs bosons was neglected; formally, the omitted terms are next-to-leading corrections in terms of the gauge couplings. The two-loop thermalization rate is thus

$$\gamma_{gb}^{(2)}/T \simeq \frac{3g^4}{256\pi} \frac{T}{m_H}. \quad (40)$$

When m_H decreases γ_{gb} increases. Therefore, for very small higgs masses the two-loop processes could actually dominate over the one-loop ones, for which the rate goes to zero as $m_H \rightarrow 0$ as is evident from Eq. (37).

There are, however, certain issues which make our two-loop estimate Eq. (40) uncertain. First, we expect $\text{Im}\Sigma(p_\mu = 0)$ to be zero (which it formally is in the full two-loop expression in [2]) but infrared dominance leads to the constant in Eq. (38), which then gives a diverging γ_H when $m_H \rightarrow 0$. Secondly, the dispersion relation with a complex Σ reads

$$\begin{aligned} \omega^2 &= p^2 + m_H^2(T) + \frac{\gamma^2}{4}, \\ \gamma &= -\frac{\text{Im}\Sigma}{\omega}, \end{aligned} \quad (41)$$

which modifies the simple relation $\omega^2 = p^2 + m_H^2(T)$ when $m_H^2(T) \lesssim |\text{Im}\Sigma|$. (Note that $\text{Re}\Sigma$ has been included in the definition of the renormalized mass m_H .) Also, when propagators have complex poles the δ -functions in Eqs. (9) and (24) should be replaced by Lorentz functions. Taken all together, we conclude that

the simple comparison with the “rising sun” diagram gives a indication that the two-loop contributions may be comparable to the one-loop ones, although the actual magnitude remains somewhat uncertain.

4. Thermalization rate vs. phase transition time

As an application, let us now turn to consider a first-order electroweak phase transition. There has been a number of attempts [1] to derive the form of the appropriate effective potential for the Higgs field in perturbation theory, but lattice studies [9] seem to indicate that there also exists a large non-perturbative component. Therefore, to be definite, we shall merely assume that the free energy of the neutral Higgs field for $\phi \simeq 0$ and near the critical temperature T_c can be expressed through the expansion

$$V(\phi) = \frac{1}{2}\kappa(T^2 - T_0^2)\phi^2 - \frac{1}{3}\alpha T\phi^3 + \frac{1}{4}\lambda\phi^4 + \dots \quad (42)$$

where κ , α , λ and T_0 are (unknown) parameters with $T_c^2 = T_0^2/(1 - 2\alpha^2/9\lambda\kappa)$. In principle, they need not be equal to the corresponding parameters at $T \simeq 0$. Thermodynamical properties of Eq. (42), as well as bubble nucleation, has been discussed in [10]. Nucleation starts at T_f where $T_0 < T_f < T_c$, and the time scale after which the major part of a given volume is filled by bubbles of new phase can be estimated to be

$$t_{\text{growth}} = -\frac{1}{S'(t_f)} = t_c \frac{\frac{1}{2}A^{\frac{1}{2}}}{\ln^{\frac{3}{2}}(M_P^4/T_c^4)} \quad , \quad (43)$$

where $S(t)$ is the action, and in the small supercooling limit $A = 0.0144 \alpha^5 \lambda^{-7/2} \kappa^{-2} (T_c/T_0)^4$. Here t_c is the time corresponding to T_c and is related to the Hubble rate by $H = (8\pi^3 g_*/90)^{1/2} T_c^2/M_P = 1/2t_c$. The size of the critical bubble, R_c , depends on the parameters of the potential Eq. (42) but is typically larger than the correlation length [11].

Thermal fluctuations generate configurations which may locally affect the dynamics of the phase transition, and for that purpose it is important to find

out the lifetime of these fluctuations. In particular, fluctuations larger than the correlation length, if sufficiently stable, could be important for bubble formation and growth.

In the previous Sections we have computed the thermalization rate for perturbations with small amplitude ($b(x)^2 \leq (p^2 + m_H^2)/\lambda$) and large spatial size ($\mathbf{p} \simeq 0$); these correspond then to large subcritical "bubbles". Since the precise T -dependence of m_H close to T_c is not reliably described by the loop expansion, we prefer to parametrize γ as a function of m_H rather than T . When varying m_H/T we can consider T to be fixed and vary the zero temperature higgs mass $m_H(0)$, which is not experimentally known. If we rather want γ as a function of T , we would have to take into account the T -dependence of m_H , and the fact that the vertical scale in Fig. 5 depends on T .

The largest fermionic contribution to the imaginary part comes from the top quark intermediate states, and in Fig. 5 we have drawn the fermionic thermalization rate, assuming $m_t = 135$ GeV and $m_L = m_R$, together with the gauge boson contributions. The thresholds are such that the one-loop rate is zero when $0.25T \lesssim m_H(T) \lesssim 1.15T$. (The threshold structure for the bottom quark is different from that of the top quark and would actually yield a non-zero thermalization rate in parts of the region $0.25T \lesssim m_H(T) \lesssim 1.15T$, but with a relatively small rate.) In contrast, the two-loop rate does not have any thresholds so that thermalization will in fact take place for all values of m_H . Comparing with Fig. 5, we find that the two-loop estimate Eq. (40) would become bigger than the one-loop rate when $m_H/T \lesssim 5 \times 10^{-2}$.

From Fig. 5 we see also that the thermalization rate is roughly $\gamma \sim 10^{-2}T \gg H$ in the region where m_H is small. Thus compared with the Hubble time, thermalization is very fast. However, the phase transition may also be very fast. This is so especially when the amount of supercooling is small. In that case typically $m_H \ll T$ during the transition and we may safely approximate $t_f \simeq t_c$. Comparing thermalization rate with the growth time we find that

$$\gamma t_{\text{growth}} \simeq 7 \times 10^{-8} A^{\frac{1}{2}} \left(\frac{M_P}{T_c} \right). \quad (44)$$

At T_c we may take for reference values the one-loop results $\kappa = 0.23$ (with $m_t = 135$ GeV) and $\alpha = 0.028$, and for λ an arbitrary value 0.006 to satisfy the baryon-number retention bound [12] (in $\lambda\phi^4$ -theories there are actually indications [13] that λ could be driven to very small values at T_c). For these values $A = 0.35$. With $T_c \simeq 100$ GeV, we then find that $\gamma t_{\text{growth}} \simeq 5 \times 10^9$. In order that $\gamma t_{\text{growth}} \sim 1$ we would need $A \sim 10^{-20}$, or extremely small α and phase transition which is almost second order. Although in principle this might be possible, we conclude that it is unlikely that spatially large fluctuations can remain stable during the electroweak phase transition.

5. Summary and conclusions

We have calculated the one-loop thermalization rate of the Higgs field in the high temperature phase in the Standard Model. A fluctuation of the Higgs field thermalizes through scattering processes with itself, with the gauge bosons and with the fermions. It is necessary to resum the hard thermal loops for the gauge boson and the fermion self-energies in order to get any kinematically allowed processes at the one-loop level. This changes the dispersion relations and induces thermal masses in such a way that the Higgs particle can annihilate with a higgs from the heat bath to create a gauge boson, be absorbed by a fermion, or decay into a pair of fermions. The inverses of these processes are also possible. By using resummed propagators for gauge bosons and fermions we consider the external states to be the effective particle excitations in the heat bath, including collective excitations.

It is the existence of the two fermionic branches at high T , particles and holes with opposite chirality-helicity relations, that allow for a Higgs particle to be absorbed by fermions in the heat bath. The unusual dispersion relation for holes makes the determination of energy thresholds complicated and it has to be done numerically. To get a positive thermalization rate one has to use propagators with the correct analytical continuation, $k_0 \rightarrow k_0 + i\epsilon k_0$, and not just add $i\epsilon$ in the

denominator. Only processes which conserve helicity are possible, and this is also essential in order to get a positive thermalization rate.

Two Higgs particles can annihilate and produce a gauge boson when the higgs mass is small enough. There are, of course, also higher order processes that do not have any thresholds. In order to make a crude estimate of the two-loop processes we estimate them by a two-loop gauge boson "rising sun" diagram. We found that for the Standard Model parameters there is a possibility that the two-loop effects are of the same order of magnitude as the one-loop ones.

The thermalization rate is compared with other time scales in Section 4. Two such scales of interest are the Hubble expansion and the electroweak phase transition rates. The time it takes the bubbles of new phase to fill up the universe is to a large extent model dependent but can be estimated in a phenomenological model [10]. We found that in general the thermalization rate is several orders of magnitude faster than both the Hubble rate and the phase transition rate. This does not mean that thermal fluctuations are unimportant for the dynamics of the phase transition. Rather, it justifies time averaging over fluctuations when comparing them with the phase transition rate.

References

- [1] For a recent review see A. G. Cohen, D. B. Kaplan and A. E. Nelson, “Progress in electroweak baryogenesis”, to appear in *Annual Review of Nuclear and Particle Science* **43**, and references therein.
- [2] P. Elmfors, K. Enqvist and I. Vilja, “Finite temperature effective action and thermalization of perturbations”, preprint NORDITA - 93/22 P.
- [3] A. Weldon, “Simple rules for discontinuities in finite-temperature field theory”, *Phys. Rev.* **D28** (1983) 2007.
- [4] R. L. Kobes and G. W. Semenoff, “Discontinuities of Green functions in field theory at finite temperature and density”, *Nucl. Phys.* **260** (1985) 714.
- [5] H. A. Weldon, “Covariant calculations at finite temperature: The relativistic plasma”, *Phys. Rev.* **D26** (1982) 1394.
- [6] V. V. Klimov, “Collective excitations in a hot quark-gluon plasma”, *Sov. Phys. JETP* **55** (1982) 199.
- [7] H. A. Weldon, “Dynamical holes in the quark-gluon plasma”, *Phys. Rev.* **D40** (1989) 2410.
- [8] R. D. Pisarski, “Renormalized fermion propagators in hot gauge theories”, *Nucl. Phys.* **A498** (1989) 423c.
- [9] K. Kajantie, K. Rummukainen and M. E. Shaposhnikov, “A lattice Monte-Carlo study of the hot electroweak phase transition”, CERN-TH.6901/93.
- [10] K. Enqvist, J. Ignatius, K. Kajantie and K. Rummukainen, “Nucleation and bubble growth in a first-order cosmological electroweak phase transition”, *Phys. Rev.* **D45** (1992) 3415.
- [11] W. Buchmüller, Z. Fodor, T. Helbig and D. Walliser, “The weak electroweak phase transition”, DESY preprint 93-021.
- [12] A. I. Bochkarev, S. V. Kuzmin and M. E. Shaposhnikov, “Model dependence of the cosmological upper bound of the Higgs-boson mass”, *Phys. Rev.* **D43** (1991) 369.

- [13] P. Elmfors, “Finite temperature renormalization for the $(\phi^3)_6$ - and $(\phi^4)_4$ -models at zero momentum”, *Z. Phys.* **C56** (1992) 601; N. Tetradis and C. Wetterich, “The high temperature phase transition for ϕ^4 -theories”, *Nucl. Phys.* **B398** (1993) 659.

Figure Captions

Fig. 1. Schematic finite temperature dispersion relations for fermions: (a) left handed particle and right handed hole; k_1 marks the maximum difference between the particle and hole energies; (b) left handed hole and right handed particle; k_2 denotes the crossing point where the energies are equal.

Fig. 2. The self-energy diagram for the fermionic thermalization rate. The arrows indicate the flow of four-momenta.

Fig. 3. The self-energy diagram for the gauge bosonic thermalization rate. The arrows indicate the flow of four-momenta.

Fig. 4. Two two-loop diagrams contributing to the thermalization rate: (a) pure scalar "rising sun"; (b) "rising sun" with gauge bosons.

Fig. 5. One-loop thermalization rates: SU(2) gauge boson contribution (dotted line), U(1) gauge boson contribution (dashed line) and top quark contribution for the absorption channel (a) and the decay channel (b) (solid lines).

Thermalization of the Higgs field at the electroweak phase transition

Per Elmfors[†], Kari Enqvist[‡] and Iiro Vilja[★]

Nordita, Blegdamsvej 17, DK-2100 Copenhagen Ø, Denmark

Abstract

The thermalization rate for long wavelength fluctuations in the Higgs field is calculated from the imaginary part of the finite temperature effective action in the unbroken phase of the Standard Model. We use improved propagators including a resummation of hard thermal loops. The thermalization rate is computed at one loop level, but an estimate of the two-loop contribution appears to give an indication that they are comparable to the one-loop result for small thermal higgs mass. We show also that the Higgs field fluctuations are likely to thermalize very fast compared with the electroweak phase transition time.

[†] internet: elmfors@nordita.dk
[‡] internet: enqvist@nbivax.nbi.dk
[★] internet: iiro@nordita.dk